

BioIn Act

PHYTOFORTIFIER
Improves plant health

Provides a significant amount of plant extracts with **PLANTEX SC®** technology

Supports by inducing the plant's natural defence system

Presents high filming capacity thanks to **ACTIVE-SET®** technology



organazoto

info@organazoto.it
www.organazoto.it



BioIn Act

Characteristics

BioIn Act is an innovative liquid activator that works synergistically to enhance the effects of BioIn Mz. **BioIn Act** is a new fertiliser designed to further increase the plant's endogenous defence mechanisms thanks to the additional plant extracts it contains, with **PLANTEX SC**® technology, soybean oil and its high content of polyunsaturated fatty acids. Among other effects, it improves the anti-transpirant and film-forming action, reduces water loss from plant tissues, improves the wetting of vegetation and reduces the activity and movement of insects with soft bodies, in which it causes dehydration. Its innovative formulation also guarantees high adhesion and distribution on treated surfaces thanks to **ACTIVE-SET**® technology.

Composition

Vegetable oils (soya oil)	100.0%
---------------------------	--------

Doses and Instructions

CROPS	FOLIAR APPLICATION	DOSE ml/hl*
Horticultural crops	During the entire growing cycle avoiding during flowering stage	200
Fruit trees	During the entire growing cycle avoiding during flowering stage	200
Cereals	After the tillering stage	200
Ornamentals	During the entire growing cycle avoiding during flowering stage	200

*The doses are indicative and should be adjusted based on the characteristics of the area and the specific needs of each crop.

Formulation	Packaging	Density (T=20°C)	pH (sol. 6 %)	Conductivity (sol. 10 %)
Soluble liquid	1-5-10-20 l	~ 980 kg/m ³	~ 6.4	~ 0.1 dS/m

Warnings

Do not exceed the maximum recommended dose of 2 L/ha per treatment. Without previous experience, it is recommended to test the product on small areas and on a limited number of plants, adjusting the dosage in case of sensitivity/phytotoxicity. Do not use in mixture with mineral oils and allow a gap between treatments with them and the application of the product. Do not apply the product near the harvest phase or during flowering. If necessary, the application can be repeated. Do not use with high temperatures (T > 30°C) or in cases of intense sunlight exposure.

Optimal results were observed when adding 50 ml/hl of BioIn MZ with 200 ml/hl of BioIn Act.

Technical Notes



organazoto

info@organazoto.it

www.organazoto.it